

NOTICE OF PUBLICATION

POND DYNAMICS/AQUACULTURE COLLABORATIVE RESEARCH SUPPORT PROGRAM



RESEARCH REPORTS

SUSTAINABLE AQUACULTURE FOR A SECURE FUTURE

Title: Bio-energetic modeling of growth and waste production of Nile tilapia (*Oreochromis niloticus* L.) in recirculation systems

Authors: M.C.J. Verdegem and A.A. van Dam
Fish Culture and Fisheries Group
Department of Animal Sciences
Wageningen University
P.O. Box 338, 6700 AH Wageningen, The Netherlands

A.A. Cabarcas-Nuñez
Department of Marine Sciences
University of Puerto Rico
P.O. Box 9013
Mayagüez, PR 00681-5000 USA

L. Oprea
Fishing and Aquaculture Department
University of Galati "Dunarea de Jos," Domneasca Str. 47
6200 Galati, Romania

Date: 15 January 2001

Publication Number: CRSP Research Report 00-162

The CRSP will not be distributing this publication. Copies may be obtained by writing to the authors.

Abstract: A bio-energetic fish growth simulation (FGS) model was developed for *Clarias gariepinus* and subsequently adjusted for the culture of *Oreochromis niloticus*, *Oncorhynchus mykiss* and *Colossoma macropomum*. The FGS model was extended with a fish waste module (FWM) to calculate the total waste production due to feeding by tilapias grown in indoor recirculation systems. Wastes calculated included the amount of uneaten feed, feces and NH_4^+ production. The amounts of wastes produced were expressed as g nitrogen (N). The model was calibrated and validated using 3 independent data sets, together comprising 175 aquarium experiments, monitoring growth in all cases and changes in proximate body composition of *O. niloticus* between stocking and harvesting in 51 cases. Fishes were grown in the individual weight range of 1 – 290 g using 32-54 % protein diets and feeding levels between 5 and 35 g $\text{kg}^{-0.8} \text{d}^{-1}$. The principal read-out parameters for calibration and validation of the model were final weight and final body fat level. Because waste production is the

Continued...

result of the same metabolic processes that lead to protein and fat deposition, it was assumed that waste production was simulated well when protein and fat deposition are. The calibrated model was used to review the effect of feeding level and dietary protein level on N-waste production per kg tilapia produced. Finally, tilapias were grown in 2 different types of recirculation systems and stagnant water ponds, quantifying N-inputs and the amount of N-wastes recovered from each system. The latter was defined as the sum of N-waste discharged (sludge and sludge water drained) and within system accumulation of N-wastes (organic and inorganic nutrients) during culture. After model calibration, the agreement between simulated and observed final weight and body fat level for all data sets was visualized. In recirculation systems different types of N-wastes were estimated well by the model. Care must be taken when applying the model to pond systems. More insight is needed on feeding ecology of tilapias in these systems.

This abstract was excerpted from the original paper, which was published in K. Fitzsimmons and J. Carvalho Filho (Editors), *Tilapia Aquaculture in the 21st Century, Fifth International Symposium on Tilapia Aquaculture*. American Tilapia Association and Departamento de Pesca e Aqüicultura/Ministério da Agricultura e do Abastecimento, Rio de Janeiro, Brazil, pp. 368–381.

CRSP RESEARCH REPORTS are published as occasional papers by the Information Management and Networking Component, Pond Dynamics/Aquaculture Collaborative Research Support Program, Oregon State University, Snell Hall 418, Corvallis, Oregon 97331-1643 USA. The Pond Dynamics/Aquaculture CRSP is supported by the US Agency for International Development under CRSP Grant No.: LAG-G-00-96-90015-00.