



PD/A CRSP NINETEENTH ANNUAL TECHNICAL REPORT

CULTURE OF MIXED-SEX NILE TILAPIA WITH PREDATORY SNAKEHEAD

*Ninth Work Plan, New Aquaculture Systems/New Species Research 2 (9NS2)
Final Report*

Yang Yi and C. Kwei Lin
Aquaculture and Aquatic Resources Management
Agricultural & Aquatic Systems and Engineering Program
Asian Institute of Technology
Pathumthani, Thailand

James S. Diana
School of Natural Resources and Environment
University of Michigan
Ann Arbor, Michigan, USA

ABSTRACT

An experiment was conducted in eighteen 200-m² fertilized earthen ponds at the Asian Institute of Technology, Thailand, from March through October 2000. This experiment was designed to assess the efficiency of snakehead (*Channa striata*) in controlling recruitment of mixed-sex Nile tilapia (*Oreochromis niloticus*) in ponds and to assess growth and production characteristics of Nile tilapia in monoculture and polyculture with snakehead. There were six treatments: A) monoculture of sex-reversed all-male tilapia; B) monoculture of mixed-sex tilapia; C) polyculture of snakehead and mixed-sex tilapia at 1:80 ratio; D) polyculture of snakehead and mixed-sex tilapia at 1:40 ratio; E) polyculture of snakehead and mixed-sex tilapia at 1:20 ratio; F) polyculture of snakehead and mixed-sex tilapia at 1:10 ratio. Sex-reversed and mixed-sex Nile tilapia were stocked at 2 fish m⁻² at sizes of 10.5 to 11.6 g and 7.2 to 8.1 g, respectively.

Results show that snakehead were able to completely control Nile tilapia recruitment at all tested predator:stocked-prey ratios, and the best predator:stocked-prey ratio was 1:80. The addition of snakehead into Nile tilapia ponds did not result in significantly greater tilapia growth, but it significantly lowered total net and gross yields of adult plus recruited tilapia. Snakehead growth was density-dependent, decreasing significantly with increasing stocking densities. While snakehead biomass gain was not significantly different at stocking densities from 0.025 to 0.1 fish m⁻², the gain was significantly lower at a stocking density of 0.2 fish m⁻². The present experiment demonstrates that snakehead are able to control Nile tilapia recruitment completely and provide an alternative technique for Nile tilapia culture.

INTRODUCTION

The aquaculture of species at lower trophic levels, such as tilapia, presents the greatest potential for efficiency (Welcomme, 1996). However, overpopulation of tilapia in confined ponds causes stunted growth due to shortage of natural food, particularly in semi-intensive culture. Various methods of population control have been applied (Mair and Little, 1991), such as culture in cages, culture with predators, intermittent harvesting, hybridization, induction of sterility, and production of super-male fish (YY-male). However, population control of tilapias by culture with predators has been practiced worldwide but not well studied. Various predatory fish species have been used with varying success in combination with different tilapia species depending on their availability. These species include snakehead (*Channa striata* or *Ophiocephalus striatus*) (Pongsuwana, 1956; Chimits, 1957; Tongsanga, 1962; Chen, 1976; Cruz and Shehadeh, 1980; Hopkins et al., 1982; Wee, 1982; Balasuriya, 1988); *Ophiocephalus obscuris* (de Graaf et al., 1996); *Micropterus salmoides* (Swingle, 1960; Meschkat, 1967; McGinty, 1985); *Lates niloticus* (Meschkat, 1967; Planquette, 1974; Lazard, 1980; Bedawi, 1985; El Gamal, 1992); *Hemichromis fasciatus* (Bardach et al., 1972; Lazard, 1980); *Cichla ocellaris* (Lovshin, 1977; McGinty, 1983; Verani et al., 1983); *Clarias* sp. (Meecham, 1975; Bard et al., 1976; Lazard, 1980; Janssen, 1985;

de Graaf et al., 1996); *Cichlasoma managuense* (Dunseth and Bayne, 1978); *Elops hawaiiensis* (Fortes, 1980); and *Megalops cyprinoides* (Fortes, 1980). However, the difficulty in breeding or obtaining predators of the correct size often resulted in limited application of this population control method (Balarin and Hatton, 1979; Penman and McAndrew, 2000).

Snakehead have long been regarded as valuable food fish and widely cultured in the Far East (Wee, 1982). It was reported to be used in polyculture with tilapia to control tilapia population or with carps to keep out other extraneous pest fish in the pond system (Wee, 1982). Snakehead are highly predacious as they swallow their prey whole (Diana et al., 1985) and have been shown to effectively prey on live tilapia fry (Kaewpaitoon, 1992). A population including 5% (predator:stocked-prey ratio of 1:20) snakehead with tilapia has been demonstrated to control tilapia recruitment (Balasuriya, 1988). Negligible tilapia recruitment was generally found during harvest where snakehead existed in tilapia ponds.

The purposes of this study were to assess:

- 1) The efficiency of snakehead in controlling overpopulation of mixed-sex Nile tilapia (*Oreochromis niloticus*) in ponds and
- 2) The growth and production of Nile tilapia in monoculture and polyculture with snakehead.

METHODS AND MATERIALS

The experiment was conducted using a randomized complete block design in eighteen 200-m² earthen ponds at the Asian Institute of Technology (AIT), Thailand. There were six treatments with triplicates, one in each block:

- A) monoculture of sex-reversed all-male tilapia;
- B) monoculture of mixed-sex tilapia;
- C) polyculture of snakehead and mixed-sex tilapia at 1:80 ratio;
- D) polyculture of snakehead and mixed-sex tilapia at 1:40 ratio;
- E) polyculture of snakehead and mixed-sex tilapia at 1:20 ratio;
- F) polyculture of snakehead and mixed-sex tilapia at 1:10 ratio.

The Chitralada strain (Thai strain) of Nile tilapia was used in the present experiment. Nile tilapia fry were obtained from the AIT Hatchery, while snakehead fingerlings were purchased from a local market. Sex-reversed Nile tilapia (10.5 to 11.6 g size) and mixed-sex Nile tilapia (7.2 to 8.1 g size) were stocked at 2 fish m⁻² in treatment A and treatments B through F, respectively, while snakehead (88.0 to 100.0 g size) were stocked at 0.025, 0.05, 0.1, and 0.2 fish m⁻² in treatments C, D, E, and F, respectively, on 30 March 2000. During the experiment, approximately 10% of the initial Nile tilapia stock was seined, counted, and weighed en masse biweekly for each pond. All

fish were harvested on 10 October 2000 after 194 days of culture. Daily weight gain (g fish⁻¹ d⁻¹), yield (kg pond⁻¹), and extrapolated yield (kg ha⁻¹ yr⁻¹) were calculated.

All ponds were dried for one month prior to the experiment to eliminate wild fish. Each pond dike was enclosed with a fine mesh net fence about 1 m tall, supported by bamboo sticks, with the lower end of the net buried in the dike soil to prevent entry of wild fish and movement of stocked snakehead from one pond to another. All ponds were fertilized with urea and triple superphosphate (TSP) at a rate of 28 kg nitrogen (N) and 7 kg phosphorus (P) ha⁻¹ wk⁻¹. Initial pond fertilization took place two weeks prior to fish stocking. Water depth in all ponds was maintained at 1 m throughout the experiment by adding water weekly to replace evaporation and seepage losses.

Integrated water samples were taken biweekly from the entire water column near the center of each pond at about 1000 h for analyses of pH, alkalinity, total ammonium nitrogen (TAN), nitrite-nitrogen, nitrate-nitrogen, total Kjeldahl nitrogen (TKN), soluble reactive phosphorus (SRP), total phosphorus (TP), chlorophyll *a*, total suspended solids (TSS), and total volatile solids (TVS) (APHA et al., 1985; Egna et al., 1989). At the time of collecting water samples, Secchi disk visibility was measured using a Secchi disk, while temperature and dissolved oxygen (DO) were measured with a YSI model 54 oxygen meter (Yellow Springs Instruments, Yellow Springs, Ohio,

Table 1. Growth performance (mean ± SE) of Nile tilapia in fertilized earthen ponds during 194-day culture. Rows of values with superscripts indicate variables with significant differences among treatments (ANOVA, $P < 0.05$). Values with similar letter superscripts are not significantly different.

Parameters	Treatments					
	A (Sex-reversed)	B (Mixed-sex)	C (1:80)	D (1:40)	E (1:20)	F (1:10)
STOCKING						
Density (fish m ⁻²)	2	2	2	2	2	2
Total Number (fish pond ⁻¹)	400	400	400	400	400	400
Mean Weight (g fish ⁻¹)	9.0 ± 0.4	7.4 ± 0.1	7.7 ± 0.1	7.5 ± 0.3	7.6 ± 0.0	7.7 ± 0.3
Total Weight (kg pond ⁻¹)	3.62 ± 0.17	2.96 ± 0.05	3.08 ± 0.04	3.01 ± 0.12	3.02 ± 0.01	3.07 ± 0.13
HARVEST						
<i>Adult Tilapia</i>						
Total Number (fish pond ⁻¹)	331 ± 10	330 ± 12	322 ± 4	312 ± 8	323 ± 9	327 ± 10
Mean Weight (g fish ⁻¹)	157.7 ± 17.5	149.8 ± 16.3	158.0 ± 10.0	152.9 ± 13.5	158.2 ± 15.3	155.3 ± 13.9
Total Weight (kg pond ⁻¹)	51.90 ± 4.60	49.03 ± 3.89	50.88 ± 3.17	47.64 ± 4.18	50.94 ± 4.05	50.60 ± 3.92
Survival Rate (%)	82.90 (78.28–87.07)	82.61 (77.97–86.81)	80.52 (75.69–84.93)	77.99 (72.97–82.63)	80.93 (76.14–85.30)	81.88 (77.17–86.15)
Daily Weight Gain (g fish ⁻¹ d ⁻¹)	0.77 ± 0.09	0.73 ± 0.08	0.77 ± 0.05	0.75 ± 0.07	0.78 ± 0.08	0.76 ± 0.07
Total Weight Gain (kg pond ⁻¹)	48.28 ± 4.43	46.07 ± 3.93	47.80 ± 3.21	44.63 ± 4.12	47.92 ± 4.06	47.53 ± 3.87
Net Yield (t ha ⁻¹ yr ⁻¹)	4.54 ± 0.42	4.33 ± 0.37	4.50 ± 0.30	4.19 ± 0.39	4.51 ± 0.38	4.47 ± 0.36
Gross Yield (t ha ⁻¹ yr ⁻¹)	4.88 ± 0.43	4.61 ± 0.37	4.79 ± 0.30	4.48 ± 0.39	4.79 ± 0.38	4.76 ± 0.37
<i>Recruited Tilapia</i>						
Total Number (fish pond ⁻¹)	---	951 ± 191	---	---	---	---
Mean Weight (g fish ⁻¹)	---	9.9 ± 0.5	---	---	---	---
Total Weight (kg pond ⁻¹)	---	9.32 ± 1.65	---	---	---	---
Net and Gross Yield (t ha ⁻¹ yr ⁻¹)	---	0.88 ± 0.15	---	---	---	---
<i>Combined Adult and Recruited Fish</i>						
Total Net Yield (t ha ⁻¹ yr ⁻¹)	4.54 ± 0.42 ^a	5.21 ± 0.22 ^b	4.50 ± 0.30 ^a	4.20 ± 0.39 ^a	4.51 ± 0.38 ^a	4.47 ± 0.36 ^a
Total Gross Yield (t ha ⁻¹ yr ⁻¹)	4.88 ± 0.43 ^a	5.49 ± 0.22 ^b	4.79 ± 0.30 ^a	4.48 ± 0.39 ^a	4.79 ± 0.38 ^a	4.76 ± 0.37 ^a

Table 2. Growth performance (mean \pm SE) of snakehead in fertilized earthen ponds during 194-day polyculture with Nile tilapia. Rows of values with superscripts indicate variables with significant differences among treatments (ANOVA, $P < 0.05$). Values with similar letter superscripts are not significantly different.

Parameters	Treatments					
	A (Sex-reversed)	B (Mixed-sex)	C (1:80)	D (1:40)	E (1:20)	F (1:10)
STOCKING						
Density (fish m ⁻²)	----	----	0.025	0.05	0.1	0.2
Total Number (fish pond ⁻¹)	----	----	5	10	20	40
Mean Weight (g fish ⁻¹)	----	----	94.5 \pm 2.4	93.7 \pm 3.5	95.8 \pm 2.2	95.7 \pm 2.2
Total Weight (kg pond ⁻¹)	----	----	0.47 \pm 0.01	0.94 \pm 0.03	1.92 \pm 0.04	3.83 \pm 0.09
HARVEST						
Total Number (fish pond ⁻¹)	----	----	4 \pm 0	8 \pm 0	19 \pm 1	35 \pm 2
Mean Weight (g fish ⁻¹)	----	----	441.3 \pm 18.1 ^a	292.0 \pm 11.6 ^b	179.3 \pm 12.0 ^c	123.3 \pm 4.2 ^d
Total Weight (kg pond ⁻¹)	----	----	1.91 \pm 0.12 ^a	2.43 \pm 0.04 ^b	3.33 \pm 0.13 ^c	4.30 \pm 0.08 ^d
Survival Rate (%)	----	----	90.75 (70.25–99.83)	83.64 (60.06–97.81)	95.47 (78.49–99.72)	88.23 (66.43–99.32)
Daily Weight Gain (g fish ⁻¹ d ⁻¹)	----	----	1.79 \pm 0.08 ^a	1.02 \pm 0.04 ^b	0.43 \pm 0.07 ^c	0.14 \pm 0.02 ^d
Total Weight Gain (kg pond ⁻¹)	----	----	1.43 \pm 0.13 ^a	1.49 \pm 0.04 ^a	1.42 \pm 0.17 ^a	0.47 \pm 0.15 ^b
Net Yield (t ha ⁻¹ yr ⁻¹)	----	----	0.14 \pm 0.01 ^a	0.14 \pm 0.00 ^a	0.13 \pm 0.02 ^a	0.05 \pm 0.01 ^b
Gross Yield (t ha ⁻¹ yr ⁻¹)	----	----	0.18 \pm 0.01 ^a	0.23 \pm 0.00 ^b	0.31 \pm 0.01 ^c	0.41 \pm 0.00 ^d

USA). Diel measurements for temperature, DO, and pH were conducted monthly in each pond at 0600, 1000, 1400, 1600, 1800, and 0600 h.

Data were analyzed statistically by analysis of variance and linear regression (Steele and Torrie, 1980) using SPSS (version 7.0) statistical software package (SPSS, Inc., Chicago, Illinois, USA). Differences were considered significant at an alpha level of 0.05. Statistical analyses for survival rates (%) were performed on the transformed data by arcsine transformation. Mean values of survival rates were given in the back-transformed scale followed by their confidence limits. All other means were given with \pm 1 standard error (SE).

A partial budget analysis was conducted to determine economic returns of the different monoculture and polyculture systems tested (Shang, 1990). The analysis was based on farm-gate prices in Thailand for harvested fish and current local market prices for all other items expressed in US dollars (US\$1 = 40 baht). Farm-gate prices of snakehead and Nile tilapia varied with size: snakehead at \$0.25 kg⁻¹ for size 100 to 200 g, \$0.50 kg⁻¹ for size 200 to 300 g, \$0.75 kg⁻¹ for size 300 to 400 g, and \$1.00 kg⁻¹ for size above 400 g; and Nile tilapia at \$0.125 kg⁻¹ for size below 50 g and \$0.375 kg⁻¹ for size 100 to 200 g. Market prices for fingerlings of snakehead (\$0.15 kg⁻¹), sex-reversed Nile tilapia (\$0.0125 piece⁻¹), and mixed-sex Nile tilapia (\$0.0042 piece⁻¹); urea (\$0.1875 kg⁻¹); and TSP (\$0.3125 kg⁻¹) were applied to the analysis. The calculation for cost of working capital was based on an annual interest rate of 8%.

RESULTS

Growth performance parameters of adult Nile tilapia were not significantly different among all treatments ($P > 0.05$, Table 1). Tilapia offspring were found only in monoculture of mixed-sex tilapia. Both sex-reversal and predator techniques were able to control the recruitment of Nile tilapia completely. However, neither sex reversal nor polyculture resulted in significantly




Figure 1. Comparison of final weight of snakehead (g) at different stocking densities (N m⁻²) of snakehead in the snakehead and Nile tilapia polyculture after 194 days.

weight; and $r = -0.985$, $P < 0.05$ for daily weight gain). Snakehead biomass gain (total weight gain and net yield) was not significantly different ($P > 0.05$) within stocking densities of 0.025 to 0.1 fish m⁻² (treatments C, D, and E; Figure 1). At a stocking density of 0.2 fish m⁻² (treatment F), snakehead biomass gain was significantly lower than in other densities ($P < 0.05$, Table 2, and Figure 2). The significantly reduced individual growth and biomass gain at the stocking density of 0.2 fish m⁻² indicated that carrying capacity of snakehead was exceeded.

The additional net yield from snakehead did not cause a significantly higher combined net yield of adult tilapia and snakehead ($P > 0.05$), but it resulted in a significantly higher

combined gross yield of adult tilapia and snakehead ($P < 0.05$, Table 3). When recruited tilapia were included, the combined net yield in the monoculture of mixed-sex tilapia (treatment B) was significantly higher than those in all other treatments ($P < 0.05$), while the combined gross yield in the monoculture of mixed-sex tilapia was similar to those in the polyculture with higher predator:stocked-prey ratios (treatments E and F, $P > 0.05$, Table 3). The results indicated that the predator:stocked-prey ratio of 1:80 (treatment C) was enough to completely control Nile tilapia recruitment.

Physical and chemical parameters of pond water were not significantly different among all treatments at all sampling times throughout the entire experimental period ($P > 0.05$). The mean values of water quality parameters were also not significantly different among all treatments ($P > 0.05$, Table 4).

The partial budget analysis (Table 5) indicated that all treatments in this experiment were profitable, and mixed-sex Nile tilapia culture (treatments B through F) produced significantly higher net return than sex-reversed Nile tilapia culture (treatment A). Snakehead and Nile tilapia polyculture at the




Figure 2. Comparison of net yield of snakehead ($\text{kg ha}^{-1} \text{yr}^{-1}$) at different stocking densities (N m^{-2}) of snakehead in the snakehead and Nile tilapia polyculture after 194 days.

Table 3. Combined yields (mean \pm SE) of Nile tilapia and snakehead in fertilized earthen ponds during 194-day culture. Rows of values with superscripts indicate variables with significant differences among treatments (ANOVA, $P < 0.05$). Values with similar letter superscripts are not significantly different.

Parameters	Treatments					
	A (Sex-reversed)	B (Mixed-sex)	C (1:80)	D (1:40)	E (1:20)	F (1:10)
ADULT TILAPIA + SNAKEHEAD						
Net Yield ($\text{t ha}^{-1} \text{yr}^{-1}$)	4.54 \pm 0.42	4.33 \pm 0.37	4.63 \pm 0.30	4.39 \pm 0.39	4.64 \pm 0.39	4.52 \pm 0.35
Gross Yield ($\text{t ha}^{-1} \text{yr}^{-1}$)	4.88 \pm 0.43 ^{abc}	4.61 \pm 0.37 ^a	4.97 \pm 0.29 ^{bc}	4.71 \pm 0.40 ^{ab}	5.11 \pm 0.38 ^c	5.16 \pm 0.36 ^c
ADULT AND RECRUITED TILAPIA + SNAKEHEAD						
Net Yield ($\text{t ha}^{-1} \text{yr}^{-1}$)	4.54 \pm 0.42 ^a	5.21 \pm 0.22 ^b	4.63 \pm 0.30 ^a	4.34 \pm 0.39 ^a	4.64 \pm 0.39 ^a	4.52 \pm 0.35 ^a
Gross Yield ($\text{t ha}^{-1} \text{yr}^{-1}$)	4.88 \pm 0.43 ^{ab}	5.49 \pm 0.22 ^c	4.97 \pm 0.29 ^{ab}	4.71 \pm 0.40 ^a	5.11 \pm 0.38 ^{bc}	5.16 \pm 0.36 ^{bc}

Table 4. Values of water quality parameters (mean \pm SE) measured throughout the experiment.

Parameters	Treatments					
	A (Sex-reversed)	B (Mixed-sex)	C (1:80)	D (1:40)	E (1:20)	F (1:10)
DO at Dawn (mg l^{-1})	2.43 \pm 0.39	2.25 \pm 0.23	2.38 \pm 0.19	2.05 \pm 0.27	2.36 \pm 0.17	2.03 \pm 0.17
Temperature ($^{\circ}\text{C}$)	27.9–37.4	27.4–36.7	27.8–37.3	27.8–36.9	27.8–36.6	27.8–35.9
pH	6.5–9.6	6.5–10.3	6.4–9.8	6.5–10.0	6.5–10.4	6.5–10.2
Alkalinity (mg l^{-1} as CaCO_3)	52 \pm 6	54 \pm 5	54 \pm 9	59 \pm 6	54 \pm 4	58 \pm 4
TKN (mg l^{-1})	6.62 \pm 0.88	6.24 \pm 0.73	6.48 \pm 0.41	5.76 \pm 0.26	6.42 \pm 0.49	6.46 \pm 0.34
TAN (mg l^{-1})	2.48 \pm 0.57	2.36 \pm 0.26	2.37 \pm 0.28	2.04 \pm 0.43	2.24 \pm 0.27	2.18 \pm 0.09
Nitrite-N (mg l^{-1})	0.19 \pm 0.04	0.22 \pm 0.01	0.20 \pm 0.01	0.14 \pm 0.02	0.21 \pm 0.04	0.21 \pm 0.02
Nitrate-N (mg l^{-1})	0.64 \pm 0.17	0.63 \pm 0.04	0.79 \pm 0.13	0.49 \pm 0.14	0.69 \pm 0.12	0.76 \pm 0.009
TP (mg l^{-1})	0.51 \pm 0.24	0.51 \pm 0.09	0.62 \pm 0.08	0.62 \pm 0.26	0.70 \pm 0.13	0.66 \pm 0.12
SRP (mg l^{-1})	0.29 \pm 0.22	0.18 \pm 0.08	0.35 \pm 0.06	0.35 \pm 0.24	0.38 \pm 0.10	0.33 \pm 0.10
Chlorophyll <i>a</i> (mg m^{-3})	35 \pm 11	46 \pm 8	31 \pm 8	42 \pm 13	44 \pm 10	41 \pm 12
Secchi Disk Visibility (cm)	24 \pm 2	24 \pm 3	25 \pm 0	22 \pm 1	24 \pm 2	23 \pm 3
TSS (mg l^{-1})	95 \pm 15	109 \pm 5	91 \pm 6	107 \pm 14	91 \pm 11	117 \pm 14
TVS (mg l^{-1})	20 \pm 3	26 \pm 2	20 \pm 1	25 \pm 4	22 \pm 4	26 \pm 4

lowest predator:stocked-prey ratio (1:80, treatment C) had the highest net return and ratio of added income to added cost, followed by the treatment at the ratio of 1:20 (treatment E).

DISCUSSION

This experiment showed that snakehead were able to completely control recruitment of Nile tilapia at a very low predator:stocked-prey ratio of 1:80, indicating high efficiency in recruitment control. A similar ratio (1:85) was used by Cruz and Shehadeh (1980) to control Nile tilapia recruitment successfully. *Lates niloticus* was reported to have a similar predation efficiency (1:84; Planquette, 1974), while other piscivorous species such as *Hemichromis fasciatus* (1:17 to 1:48; Bardach et al, 1972; Lazard, 1980), *Cichla ocellaris* (1:15; Lovshin, 1977), *Clarias lazera* (1:10; Bard et al., 1976), *Cichlasoma managuense* (1:4; Dunseth and Bayne, 1978), *Elops hawaiiensis* (1:20; Fortes, 1980), *Megalops cyprinoides* (1:10; Fortes, 1980), *Clarias gariepinus* (1:2.7; de Graaf et al., 1996), and *Ophiocephalus obscuris* (1:30; de Graaf et al., 1996) were less effective.

The present experiment clearly showed that the carrying capacity of snakehead was exceeded at the predator:stocked-prey ratio of 1:10, and poor growth occurred due mainly to the limited food items available. Snakehead is carnivorous and highly predacious on aquatic organisms such as insects; fish, including its own species; frogs; shrimps; and even small aquatic snakes (Wee, 1982). In another study snakehead had better growth at the same predator:stocked-prey ratio with smaller stocking size (0.3 g) and harvest size (108.5 g; Balasuriya, 1988). In comparison, stocking size was 95.7 g in the present experiment. The results of the present experiment suggest that the standing crop of snakehead at a stocking density of 2 Nile tilapia m⁻² should be below 4.30 kg per 200 m², or 215 kg ha⁻¹, to achieve good growth.

In other studies, high yields of harvestable-size tilapia were reported, and the final size of harvested tilapia increased with effective predators (Swingle, 1960; Lovshin, 1977; Dunseth and Bayne, 1978; Edwards et al., 1994). However, there were no significant differences in final size and yield of harvested adult tilapia among treatments in this study, and total production combining adult and recruited tilapia was significantly reduced in all polyculture treatments as the recruits were eaten. This is consistent with the results using other piscivorous species reported by Maar et al. (1966), Lovshin (1977), Fortes (1980), McGinty (1983, 1985), and Edwards et al. (1994).

The growth of sex-reversed all-male Nile tilapia was only 5% faster than mixed-sex tilapia, and this difference was not statistically significant in the present experiment. In comparison, the sex-reversed all-male tilapia grew more than 10% faster than mixed-sex tilapia in other experiments (Pascual and Mair, 1997). Stunting with mixed-sex tilapia culture, caused by competition for food between recruits and stocked tilapia, was not observed in the present experiment. Green and Teichert-Coddington (1994) also did not find significant differences between sex-reversed and mixed-sex Nile tilapia growth in ponds. Dan and Little (2000) reported that growth difference between sex-reversed and mixed-sex Thai strain of Nile tilapia (new-season seed) was significant when cultured in ponds but not significant when cultured in cages. Clearly, there are system-specific differences that may affect the growth, production, and stunting of mixed-sex tilapia.

ANTICIPATED BENEFITS

The results of this study demonstrated that snakehead can control Nile tilapia recruitment completely at low predator:stocked-prey ratios. This provides an alternative technique for Nile tilapia culture system. Tilapia recruitment control by stocking predators can make Nile tilapia production expand

Table 5. Partial budget analysis (US\$) for sex-reversed and mixed-sex Nile tilapia monoculture (treatments A and B) and snakehead and mixed-sex Nile tilapia polyculture (treatments C through F) in the 194-day experiment (based on 200-m² ponds).

Parameters	Treatments					
	A (Sex-reversed)	B (Mixed-sex)	C (1:80)	D (1:40)	E (1:20)	F (1:10)
GROSS REVENUE						
Adult Tilapia	19.46	18.39	19.08	17.87	19.10	18.98
Recruited Tilapia	----	1.16	----	----	----	----
Snakehead	----	----	1.91	1.82	1.67	1.08
Total	19.46	19.55	20.99	19.69	20.77	20.05
VARIABLE COST						
Tilapia Fingerlings	5.00	1.67	1.67	1.67	1.67	1.67
Snakehead Fingerlings	----	----	0.07	0.14	0.29	0.58
Urea	6.75	6.75	6.75	6.75	6.75	6.75
TSP	6.56	6.56	6.56	6.56	6.56	6.56
Cost of Working Capital	0.78	0.64	0.64	0.64	0.65	0.66
Total cost	19.09	15.62	15.69	15.76	15.92	16.22
NET RETURN	0.37	3.93	5.30	3.92	4.85	3.83
ADDED COST	3.33	----	0.07	0.14	0.29	0.57
ADDED RETURN	-0.09	----	1.44	0.14	1.22	0.50
ADDED INCOME / ADDED COST	-0.03	----	20.41	0.97	4.22	0.87

and increase, especially in rural areas where sex-reversed tilapia are not available. It will benefit culturists throughout Southeast Asia and other regions where tilapia are commonly cultured, snakehead are available, and there is no tilapia hatchery for sex-reversed all-male fry.

ACKNOWLEDGMENTS

The authors wish to acknowledge the Asian Institute of Technology, Thailand, for providing the research, field, and laboratory facilities. Mr. Chumpol S., Mr. Manoj Y., and Mr. Supat P. are greatly appreciated for their field and laboratory assistance.

LITERATURE CITED

- APHA, 1985. Standard Methods for the Examination of Water and Wastewater, Sixteenth Edition. American Public Health Association, American Water Works Association, and Water Pollution Control Federation, Washington, DC, 1,268 pp.
- Balarin, J.D. and J.P. Hatton, 1979. Tilapia. A guide to their biology and culture in Africa. Unit of Aquatic Pathobiology, University of Stirling, Scotland, 174 pp.
- Balasuriya, C., 1988. Snakehead (*Channa striata*) as a controlling predator in the culture of Nile tilapia (*Oreochromis niloticus*). M.Sc. thesis, Asian Institute of Technology, Bangkok, Thailand.
- Bard, J., P. de Kimpe, J. Lazard, J. Lemasson, and P. Lessent, 1976. Manuel de Pisciculture Tropicale. Centre Technique Forestier Tropical, Nogent-sur-Marine, France, 209 pp. (in French)
- Bardach, J.E., J.H. Ryther, and W.O. McLarney, 1972. Aquaculture, the Farming and Husbandry of Freshwater and Marine Organisms. Wiley-Interscience Inc., New York, 868 pp.
- Bedawi, R.M., 1985. Recruitment control and production of market size *Oreochromis niloticus* with the predator *Lates niloticus* L. in the Sudan. *J. Fish Bio.*, 26:459-464.
- Chen, T.P., 1976. Aquaculture Practices in Taiwan. Fishing News Books Ltd., Farnham, England, 162 pp.
- Chimits, P., 1957. The *Tilapia* and their culture, a second review and bibliography. *FAO Fish. Bull.*, 10(1):1-24.
- Cruz, E.M. and Z.H. Shehadeh, 1980. Preliminary results of integrated pig-fish and duck-fish production tests. In: R.S.V. Pullin and Z.H. Shehadeh (Editors), Integrated Agriculture-Aquaculture Farming Systems. ICLARM Conf. Proc. 4. International Center for Living Aquatic Resources Management, Manila, and the Southeast Asian Regional Centre for Graduate Study and Research in Agriculture, Laguna, Philippines, pp. 235-238.
- Dan, N.C. and D.C. Little, 2000. The culture performance of monosex and mixed-sex new-season and overwintered fry in three strains of Nile tilapia (*Oreochromis niloticus*) in northern Vietnam. *Aquaculture*, 184:221-231.
- de Graaf, G.D., F. Galemoni, and B. Banzoussi, 1996. Recruitment control of Nile tilapia, *Oreochromis niloticus*, by the African catfish, *Clarias gariepinus* (Burchell 1822), and the African snakehead, *Ophiocephalus obscuris*. 1. A biological analysis. *Aquaculture*, 146:85-100.
- Diana, J.S., W.Y.B. Chang, D.R. Ottey, and W. Chuapohuk, 1985. Production systems for commonly cultured freshwater fishes of southeast Asia. International Programs Report, No. 7. Great Lakes and Marine Waters Center, University of Michigan, Ann Arbor, Michigan.
- Dunseth, D.R. and D.R. Bayne, 1978. Recruitment control and production of *Tilapia aurea* (Steindachner) with the predator *Cichlosoma managuense* (Günther). *Aquaculture*, 14:383-390.
- Edwards, P., C. Pacharaprakiti, and M. Yomjinda, 1994. An assessment of the role of buffalo manure for pond culture of tilapia. 1. On-station experiment. *Aquaculture*, 126:83-95.
- Egna, H.S., N. Brown, and M. Leslie, 1989. Pond Dynamics/Aquaculture Collaborative Research Data Reports, Volume 1, General Reference: Site Descriptions, Materials and Methods for the Global Experiment. Pond Dynamics/Aquaculture CRSP, Oregon State University, Corvallis, Oregon, 84 pp.
- El Gamal, A.A., 1992. Predation by Nile perch *Lates niloticus* (L.) on *Oreochromis niloticus* (L.), *Cyprinus carpio* (L.), *Mugil* sp. and its role in controlling tilapia recruitment in Egypt. *J. Fish Biol.*, 40:351-358.
- Fortes, R.D., 1980. Tarpon as predator to control Java tilapia young in brackishwater ponds. *Fish Res. J. Philipp.*, 5(2):22-35.
- Green, B.W. and D.R. Teichert-Coddington, 1994. Growth of control and androgen-treated Nile tilapia during treatment, nursery and grow-out phases in tropical fish ponds. *Aquacult. Fish. Manage.*, 25:613-621.
- Hopkins, D.K., D. Pauly, E.M. Cruz, and J.M. van Weerd, 1982. An alternative to predator-prey ratios in predicting recruitment. *Meeresforschung Rep. Mar. Res.*, 29(3): 125-134.
- Janssen, J., 1985. Elevage du poisson-chat africain *Clarias lazera* (Cuv. and Val., 1840) en Republique Centrafricaine. 3. Alevinage et grossissement en etangs. Document Technique No. 22. FED-DGIS/FAO, Bangui, Central African Republic, 32 pp. (in French).
- Kaewpaitoon, K., 1992. Utilization of septage-raised tilapia (*Oreochromis niloticus*) as a feed for snakehead (*Channa striata*). Ph.D. dissertation, Asian Institute of Technology, Bangkok, Thailand.
- Lazard, J., 1980. Le developpement de la pisciculture intensive en Côte-d'Ivoire. Exemple de la ferme piscicole pilote de Natio-Kobadara (Korhogo). *Bois Fôrets Trop.*, 190:45-66.
- Lovshin, L.L., 1977. Progress Report on Fisheries Development in Northeast Brazil. Research and Development Series, No. 14. International Center for Aquaculture, Agricultural Experiment Station, Auburn University, Auburn, Alabama, 11 pp.
- Maar, A., M.A.E. Mortimer, and I. Van der Lingen, 1966. Fish Culture in Central East Africa. Food and Agriculture Organization, Rome, Italy, 158 pp.
- Mair, G.C. and D.C. Little, 1991. Population control in farmed tilapias. *Naga*, 14(3):8-13.
- McGinty, A.S., 1983. Population dynamics of peacock bass, *Cichla ocellaris* and *Tilapia nilotica* in fertilised ponds. In: L. Fishelson and Z. Yaron (Editors), Proc. Int. Symp. on Tilapia Aquaculture, Tel Aviv University, Nazareth, Israel, pp. 86-94.
- McGinty, A.S., 1985. Effects of predation by largemouth bass in fish production ponds stocked with *Tilapia nilotica*. *Aquaculture*, 46:269-274.
- Meecham, K., 1975. Aquaculture in Malawi. Paper presented at the FAO/CIFA Symposium on Aquaculture in Africa. Accra, Ghana. CIFA/75/SC1, FAO, Rome.
- Meschkat, A., 1967. The status of warm-water fish culture in Africa. *FAO Fish. Rep.*, 44(2):88-122.
- Pascual, L.P. and G.C. Mair, 1997. Culture performance of genetically male tilapia (GMT). In: G.C. Mair and T.A. Abella (Editors), Technoguide on the Production of Genetically Male Tilapia (GMT). Freshwater Aquaculture Center, Central Luzon State University, Nueva Ecija, Philippines, pp. 27-38.
- Penman, D.J. and B.J. McAndrew, 2000. Genetics for the management and improvement of cultured tilapias. In: M.C.M. Beveridge and B.J. McAndrew (Editors), *Tilapia: Biology and Exploitation*. Kluwer Academic Publishers, The Netherlands, pp. 227-266.
- Planquette, P., 1974. Nile perch as a predator in tilapia ponds. *FAO Aquacult. Bull.*, 6:7.
- Pongsuwana, U., 1956. Production of *Tilapia mossambica* in an experimental pond at Bangkok, Thailand. *IPFC Proc.*, 6(2):197-201.
- Shang, Y.C., 1990. Aquaculture Economic Analysis: An Introduction. World Aquaculture Society, Baton Rouge, Louisiana, 211 pp.
- Steele, R.G.D. and J.H. Torrie, 1980. Principles and Procedures of Statistics: A Biometrical Approach, Second Edition. McGraw-Hill, New York, 633 pp.
- Swingle, H.S., 1960. Comparative evaluation of two tilapias as pondfishes in Alabama. *Trans. Amer. Fish. Soc.*, 89:142-148.
- Tongsanga, S., 1962. A preliminary report on the combination of plachon (*Ophiocephalus striatus* Bloch) and tilapia (*Tilapia mossambica* Peters) in Thailand. *IPFC Proc.*, 10(2):174-180.
- Verani, J.R., M.A. Marins, A.B. da Silva, and A.C. Sobrinho, 1983. Population control in intensive fish culture associating *Oreochromis niloticus* with the natural predator *Cichla ocellaris*, a quantitative analysis. In: L. Fishelson and Z. Yaron (Editors), Proc. Int. Symp. on Tilapia Aquaculture. Tel Aviv University, Nazareth, Israel, pp. 580-587.

- Wee, K.L., 1982. The biology and culture of snakeheads. In: J. Muir and R.J. Roberts (Editors), *Recent Advances in Aquaculture*. Croom Helm Press, London, pp. 179–213.
- Welcomme, R.L., 1996. Aquaculture and world aquatic resources. In: D.J. Baird, M.C.M. Beveridge, L.A. Kelly, and J.F. Muir (Editors), *Aquaculture and Water Resource Management*. Blackwell Science, Ltd., London, pp. 1–18.

